

Signs of an adrenal crisis

Symptoms can vary in individuals but the following can be signs of an impending crisis:-

- * Severe weakness
- * Dizziness
- * Floppiness
- * Failure to respond
- * Nausea and vomiting
- * Pallor and clammy sweating
- * Severe sweating with weakness
- * Cold hands and feet
- * Pins and needles or numbness in hands or feet
- * Tachycardia (fast heart beat)
- * Hypotension (low blood pressure)
- * Hypoglycaemia (low blood sugar)

Glucogel for low blood sugar

- * Glucogel® (used to be known as HypoStop) should be only used when your child is showing signs of low blood sugar levels.
- * Glucogel® should be used AFTER your child has had the hydrocortisone injection.

How to administer Glucogel


Gradually squirt the Glucogel® into the side of your child's mouth, between the gums and the cheek. Alternatively, squirt the Glucogel® onto your fingertip and apply it between your child's gums and cheek. Up to one-third of a 25g tube may be needed. Massage your child's cheek to allow the gel to be absorbed, this should raise your child's blood sugar level within 10 minutes.

Important points

- * The hydrocortisone injection will only last in the body, for 4-6 hours, depending on how long the body takes to use it up.
- * It works quicker (peaks within approx. 20 minutes as opposed to approx. 2 hours for tablets) as it is administered directly into the muscle so it does not need to go through the stomach

Note that a higher dose of hydrocortisone does not last longer in the blood stream than a lower dose, so it is not appropriate to give more than the recommended dose. The reason it does not last longer is a bit complicated but it has to do with how the body removes medicines from the body and in fact adding more in, will cause the body to get rid of it quicker. If too high a dose is given the body's natural response will be to excrete the excess as soon as possible.

The graph below shows the differences in the cortisol when measured in the blood after the same amount of hydrocortisone is given orally and IV.



If your child should require emergency surgery a bolus of hydrocortisone must be given before surgery and protocol followed regarding hydrocortisone dosing during or after surgery. Please contact the Endocrine Registrar on call for advice on the protocol

Treating Diarrhoea

Diarrhoea is a particular problem due to the fluid losses. Again double/triple the dose of hydrocortisone is needed. Oral rehydration solutions such as Dioralyte should be used. It is important to make sure that your child is passing urine regularly. Fludrocortisone dose must not be increased.

Seek medical advice early especially if there is:-

- * a fever
- * blood in the diarrhoea
- * the child becomes confused
- * the diarrhoea does not stop after 24 hours.

Do not use anti diarrhoeal drugs.

ALWAYS REMEMBER: If in doubt it is safer to give the injection than not to give it, as it will do no real harm if given as a one off emergency dose, even if it is not really needed.

Professor Peter Hindmarsh
Addison's Clinic
Great Ormond Street Hospital for Children
NHS Foundation Trust
London WC1N 3JH
Phone: 020 7405 9200

NHS

Great Ormond Street
Hospital for Children

NHS Foundation Trust

Adrenal
Insufficiency
Addison's

ADDISON'S

A GUIDE FOR ILLNESS
AND
EMERGENCIES

Professor Peter Hindmarsh

The Endocrine Team at Great Ormond Street provides 24 hour cover every day of the year. For urgent advice phone 0207 405 9200 and ask for the Medical Registrar on call who is always backed up by on-call Endocrine Consultant

WHAT TO DO IN ILLNESS AND INJURIES

We realise that when first faced with the diagnosis of your child having Addison's one of the most daunting and worrying issues is dealing with illness and emergencies, especially in babies and young toddlers who are not able to communicate on how they feel. We hope the following guidelines will help.

So how do we deal with Emergencies and Illness?

Remember that you know your child best but here are some steps, which we have put into Levels of illness that you can follow.

Level 1

- * Your child is unwell and has a temperature over 38°C.
- * He/she is taking fluids.

In this situation then the hydrocortisone dose should be doubled or trebled. If you are using twice a day hydrocortisone you may well need to increase the frequency of dosing to three times per day. This would mean adding in an extra dose at lunchtime and the amount would be double the dose that you use in the morning. To cover the early hours of the morning when the night time dose has worn off, we now advise an extra double dose to be given around 4 am, this is to ensure there is cortisol around and will also go some way to preventing low blood glucose.

Summary

- * Morning double/triple dose
- * Lunchtime double/triple morning dose
- * Bedtime double/triple dose
- * We also recommend an extra double morning dose at 4am

Fluids that contain some sugar should be encouraged.

Level 2

- * This would be same situation as in Level 1 but there is also associated vomiting.
- * If your child has vomited 2-4 hours after the last dose of hydrocortisone then the last dose has probably been absorbed. But if he/she vomits within 2 hours of the last dose then
- * Give repeat double or triple dose orally. If this stays down, continue with double or triple dose orally.
- * If not and your child vomits it back within 2 hours then intramuscular hydrocortisone should be administered.

Intramuscular Hydrocortisone Doses For Emergency Use

Dose (mg)	Age range (years)
25	0 - 1
50	1 - 5
100	over 5

Following the administration of the intramuscular injection of hydrocortisone take your child to the nearest Accident and Emergency department as soon as it is possible and give detailed explanation to the nurses and doctors that emergency hydrocortisone has already been administered. Show them your emergency letter and Adrenal Insufficiency Card.

You need to make sure that the medical team measure the following and you should not leave until you have the results:

- * Urea and electrolyte concentrations to make sure salt levels are not low
- * Blood glucose concentration as this can go low
- * Blood Pressure which if low tells us if adrenal crisis is pending

Should any of these levels be abnormal or in any situation of doubt then your child should be admitted for glucose, electrolyte and blood pressure monitoring along with possible need for intravenous fluids and hydrocortisone.

We recommend that after a hydrocortisone injection is given, a child should be admitted for observation for at least 12 hours.

The hydrocortisone injection should give cover for around 4 - 6 hours, so double or triple dose needs to be resumed after this time which will depend on how well your child has recovered, i.e.: if your child starts to show signs of getting worse again after 4 hours, triple dose should be given.

Injuries and Accidents

- * For mild bumps, scratches and bruises, there is no need to increase the dose
- * For sprains and other painful injuries, it will do no harm to give a double dose
- * For burns and serious injuries such as broken bones, head injury, severe fall with the possibility of a broken bone - give the intramuscular hydrocortisone injection.

If your child is unconscious, give the hydrocortisone injection first, then dial 999 and state that your child has 'Adrenal Insufficiency' See the section on Calling an ambulance for important information.

Why do we use the Intramuscular Injection?

It does sound drastic but like any child once they start vomiting it is hard for them to keep down fluids and in your child's case hydrocortisone. Also whatever makes them vomit may also interfere with the absorption of hydrocortisone from the stomach so the intramuscular injection bypasses this.

When to Call Ambulance:

You should call an ambulance when:

1. **AFTER you have given an emergency injection** and there has been no response
2. If your child is involved in an accident
3. There is loss of consciousness
4. Your child is vomiting and cannot keep medicines down
5. If your child is very unwell and you do not have an injection kit or unable to give the injection.

Calling an Ambulance:-

Many ambulance services in the UK are now following a 'pathways system' and each 999 call is triaged.

The 'pathway' followed for Addison's and every other cortisol deficient disorder is -

'ADRENAL INSUFFICIENCY.'

The call handler will then ask for various details such as name, date of birth and what is wrong - it is most important that you state that the patient has **ADRENAL INSUFFICIENCY and NEEDS AN EMERGENCY HYDROCORTISONE INJECTION.**

ADRENAL INSUFFICIENCY is the key statement for the Ambulance Crew who will recognise it, respond accordingly and will act by giving emergency hydrocortisone

At present if you have a protocol logged with emergency services this is flagged by the patient's address (not by their name) so it is important to make the emergency call handler aware that there is a protocol in place and to give the patient's address if you have it.

When the ambulance arrives give a copy of our emergency letter or Adrenal Insufficiency Cards to the team.

Note that if an extra dose of hydrocortisone is given, this will not be harmful in any way.

On the Way to Hospital:

Give the Paediatric Endocrine team at Great Ormond Street Hospital for Children London a call (24 hour cover) as soon as possible on 020 7405 9200 and ask to be put through to the Paediatric Registrar on call.

Medic Alert Please note this reinforces the importance of wearing a medic alert with the wording **ADRENAL INSUFFICIENCY** and also carrying one of our Adrenal Insufficiency Cards which will have the home address details on.